



HERMIT PARK VET HOSPITAL
 223 Charters Towers Road
 Hermit Park QLD 4812
 P: 4779 5222 F: 4779 9239

BAYSIDE VET CLINIC
 34 Primrose Street
 Belgian Gardens QLD 4810
 P: 4772 1513 F: 4724 1957

Dr Grace Leong B.V.Sc., B.Sc.
 Dr Kimberley Nagle B.V.Sc.
 & Associates
 E: palmsvetgroup@palmsvetgroup.com.au
 www.palmsvetgroup.com.au

TICK PREVENTATIVES AVAILABLE FOR DOGS

APPLICATION TYPE	PRODUCT NAME	INFORMATION
Oral Chew	NexGard	Monthly flavoured chew for fleas and ticks (bush, brown dog and paralysis ticks).
	Bravecto	Flavoured chew for dogs that protects against paralysis tick for 4 months, fleas 3 months and brown dog ticks for 2 months. We recommend you give the product every 3mths unless you are having a problem with brown dog ticks in which case we recommend every 2 mths.
Top-Spot	Advantix	Applied to back of the neck. Lasts 1mth against fleas and brown dog ticks but needs to be applied fortnightly when travelling in paralysis tick areas. This product also repels ticks so they are less likely to attach. It also repels other biting parasites such as mosquitos and sandflies. TOXIC TO CATS!
	Frontline Plus	Use monthly against Brown dog ticks and fleas. Apply fortnightly in paralysis tick areas.
Tick Collars	Scalibor	Protects against paralysis ticks 14 weeks and brown dog ticks 26 weeks. Needs to be applied at least 2-3weeks before exposure to ticks. Remove for swimming/ bathing.
	Kiltix	Flea and tick collar. Controls fleas, brown dog tick and bush ticks for 5mths and paralysis ticks for 6weeks. Remove for swimming/bathing.
Leave on Concentrates	Permaxin	Leave on rinse applied weekly. Reapply after swimming/bathing. TOXIC TO CATS
	Fidos Flea and Tick Concentrate	Applied as a leave on rinse every 3 days. Needs reapplication after swimming/bathing